User Guide VaPro Pocket[™] Catheters















5. Remove the catheter from the package. Leave pack on the lap, sink or flat surface. Lay catheter on opened package.







- 7. Remove the cap from the protective tip of catheter.
- 8. Hold the catheter in one hand, and with the other hand advance the catheter forward until tip of the catheter fills the protective tip, taking care that the catheter does not protrude from the tip.











- 10. Using your fingers and forefinger of your non-dominant hand squeeze penis gently to open the urethral opening. With your dominant hand, insert the protective tip until the plate comes in contact with urethral opening.
- 11. Insert the catheter forward until urine starts to flow.





- 12. Once urine flow has stopped slowly withdraw the catheter. If urine flows again, pause until urine stops. After you're sure the bladder is empty, remove the catheter completely. Reposition the foreskin (if present).
- Dispose of the catheter and packaging in a disposal bin. DO NOT FLUSH DOWN TOILET.
- 14

14. Wash your hands.

WARNING: To help reduce the potential for infection and/or other complications, do not reuse.

WARNING: If discomfort or any sign of trauma occurs, discontinue use immediately and consult your healthcare practitioner.

- Ready-to-use, single-use intermittent catheter
- Protective tip and sleeve supports no-touch insertion technique
- · Colour-coded funnel
- . Not made with natural rubber latex

- Hydrophilic-coated, Phthalate-free catheter
- 2 smooth catheter eyelets
- · Easy-to-remove ring cap

INDICATIONS FOR USE

This intermittent catheter is a flexible tubular device that is inserted through the urethra by male, female and paediatric patients who need to drain urine from the bladder.

PRECAUTION: Please consult your healthcare practitioner before using this product if any of the following conditions are present:

- · Severed urethra
- · Pronounced stricture
- Urethritis inflammation of the urethra
- Epididymitis inflammation of the epididymis (testicle tube)
- Unexplained urethral bleeding
- False passage
- Prostatitis inflammation of the prostate gland

PRECAUTION: Self-catheterisation should follow the plan of care and advice given by your healthcare practitioner and be carried out only in accordance with the instructions for use provided. Because catheterisation frequency varies by person, the recommended frequency of your catheterisation should be provided by your healthcare practitioner. For any other questions about your catheterisation, please contact your healthcare practitioner.

NOTE: Store boxes in a flat position and at normal room temperature. This product consists of: polyurethane sleeve, thermoplastic elastomer tip, and tubing not made with DEHP.

Rx only

Hollister Limited

Building 1010, Winnersh Triangle Business Park, Eskdale Road, Winnersh, Wokingham RG41 5TS

For more information and to request FREE samples:

Tel: 0800 521 377 | Email: samples.uk@hollister.com

www.hollister.co.uk

Scan to watch video



